





S NCBI	Pubywea	of Medicine NLM		
PubMed Nucleotide Search PubMed ▼ for	The state of the s	PopSet	Taxonomy Go Cl	OMIM ear
Scalor and a second	The state of the s	tory 🤚	Clipboard	5
Entrez PubMed	Display Abstract Save Text.	Order	Add to Clipbo	ard
	1: Zentralbl Bakteriol 1995 Oct;282(4):4	116-26	Related Art	ticles, Books
PubMed Services Immunogenicity and antigenic relationship of Salmonella enterotoxin with other enterotoxins.				
	Rahman H, Sharma VD.			
	Department of Microbiology and Public Health, G. B. Pant University of Agriculture and Technology, Pantnagar, Nainital, India.			
Related Resources	The immunogenicity and antigenic relationship of Salmonella enterotoxin with other enterotoxins were studied. The purified enterotoxin of Salmonella typhimurium strains was immunogenic in rabbits and antiserum produced against it completely neutralized the enterotoxic activity of purified as well as crude enterotoxins. The Salmonella enterotoxin did not cross-react with cholera toxin, heat-labile enterotoxin of Escherichia coli or Shiga toxin. On the other hand, enterotoxins produced by different serotypes of Salmonella were antigenically related. Heat, pH and proteolytic enzymes had not only			

were antigenically related. Heat, pH and proteolytic enzymes had not only adversely affected the enterotoxigenicity but also the antigenicity of the Salmonella enterotoxin. However, formalin was found to completely destroy the enterotoxigenicity of enterotoxin without affecting the antigenicity. Formalin-treated enterotoxin may thus be tried as a vaccine to prevent the diarrhoeal syndrome induced by enterotoxigenic salmonellae.

PMID: 9810665 [PubMed - indexed for MEDLINE]

Text : Order : Add to Clipboard Display : Abstract

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer